arXiv:0903.3835v1 [math.AG] 23 Mar 2009

Poincaré series of collections of plane valuations *

A. Campillo F. Delgado S.M. Gusein-Zade
F. Hernando †

Abstract

In earlier papers there were given formulae for the Poincaré series of multi-index filtrations on the ring $\mathcal{O}_{\mathbb{C}^2,0}$ of germs of functions of two variables defined by collections of valuations corresponding to (reducible) plane curve singularities and by collections of divisorial ones. It was shown that the Poincaré series of a collection of divisorial valuations determines the topology of the collection of divisors. Here we give a formula for the Poincaré series of a general collection of valuations on the ring $\mathcal{O}_{\mathbb{C}^2,0}$ centred at the origin and prove a generalization of the statement that the Poincaré series determines the topology of the collection.

In [1], [6], ... there were considered multi-index filtrations on the ring $\mathcal{O}_{\mathbb{C}^2,0}$ of germs of functions of two variables defined by collections of valuations corresponding to (reducible) plane curve singularities and by collections of divisorial ones. One gave formulae for the Poincaré series of such filtrations in terms of an embedded resolution of the curve singularity or of the collections of divisors respectively. These formulae give Poincaré series as rational functions equal to products/ratios of cyclotomic polynomials. In particular, it was shown that the Poincaré series of the collection of valuations corresponding to a curve coincides with the Alexander polynomial of the corresponding algebraic link. This implies that the Poincaré series of such a collection determines the topology of the curve ([7]). An analogue of this statement for divisorial valuations was proved in [3]. Here we give a formula for the Poincaré series of a general collection of valuations on the ring $\mathcal{O}_{\mathbb{C}^2,0}$ centred at the origin. We also prove a generalization of the statement that the Poincaré series determines the topology of the collection of divisors.

---

*Math. Subject Class.: 14H20, 32S45, 13F30. Key words: Poincaré series, plane valuations. Partially supported by the grant MTM2007-64704. Third author is also partially supported by the grants RFBR-007-00593 and NSh-709.2008.1.

†Addresses: A. Campillo and F. Delgado: University of Valladolid, Dept. of Algebra, Geometry and Topology, 47011 Valladolid, Spain. E-mail: campillo@agt.uva.es, fdelgado@agt.uva.es; S.M.Gusein-Zade: Moscow State University, Faculty of Mathematics and Mechanics, Moscow, GSP-1, 119991, Russia. E-mail: sabir@mccme.ru; F. Hernando: Dept. of Mathematics. University College Cork. Ireland. E-mail: f.hernando@ucc.ie

1 Poincaré series of several valuations

Let $\mathcal{O}_{X,0}$ be the ring of germs of functions on a germ $(X, 0)$ of a complex analytic variety and let G be a ordered abelian group, $G^+ := \{a \in G : a \geq 0\}$. A valuation on the ring $\mathcal{O}_{X,0}$ with values in the group G is a map $v : \mathcal{O}_{X,0} \to G^+ \cup \{+\infty\}$ such that:

1. $v(g_1 \cdot g_2) = v(g_1) + v(g_2)$;
2. $v(g_1 + g_2) \geq \min\{v(g_1), v(g_2)\}$;
3. $v(\lambda) = 0$ for $\lambda \in \mathbb{C}^* = \mathbb{C} \setminus \{0\}$.

The semigroup $S = \operatorname{Im} v \setminus \{+\infty\}$ (the semigroup of values of the valuation v) is well ordered (i.e. each subset of S has the minimal element) and moreover each element $a \in S$ has a finite number of representations as the sum $a_1 + a_2$ of two elements of S ([8], see also [4]).

Let v_i, $i = 1, \ldots, s$, be valuations on the ring $\mathcal{O}_{X,0}$ with values in ordered groups G_i and the semigroups of values S_i. The direct product $S = S_1 \times \ldots \times S_s$ is a partially ordered semigroup: $\underline{v}' = (v'_1, \ldots, v'_s) \geq \underline{v}'' = (v''_1, \ldots, v''_s)$ iff $v'_i \geq v''_i$ for all $i = 1, \ldots, s$ ($v'_i, v''_i \in S_i$). For a germ $g \in \mathcal{O}_{X,0}$, let $\underline{v}(g) := (v_1(g), \ldots, v_s(g))$.

Definition: The ring $\mathbb{Z}[[S]]$ of power series on the semigroup $S = S_1 \times \ldots \times S_s$ is the set of formal expressions of the form $\sum\limits_{\underline{v} \in S} a_{\underline{v}}\, \underline{t}^{\underline{v}}$ ($\underline{v} := (v_1, \ldots, v_s) \in S$, $\underline{t}^{\underline{v}} := t_1^{v_1} \cdot \ldots \cdot t_s^{v_s}$) with the usual ring operations.

Remarks. **1.** The fact that $\mathbb{Z}[[S]]$ is a ring (i.e. that the multiplication is defined) follows from the described properties of the semigroup of values of a valuation.

2. If $G_i = \mathbb{Z}$ for $i = 1, \ldots, s$, the ring $\mathbb{Z}[[S]]$ is contained in the ring $\mathbb{Z}[[t_1, \ldots, t_s]]$ of power series in the variables $t_1, \ldots, t_s$ with integer coefficients and thus an element of the ring $\mathbb{Z}[[S]]$ is a usual power series in the variables $t_1, \ldots, t_s$.

The collection $\{v_i\}$ of valuations defines a multi-index filtration on the ring $\mathcal{O}_{X,0}$ by the ideals

$$J(\underline{v}) = \{g \in \mathcal{O}_{X,0} : \underline{v}(g) \geq \underline{v}\}$$

(indexed by the elements $\underline{v}$ of the group $G = G_1 \times \ldots \times G_s$). For $I \subset I_0 = \{1, \ldots, s\}$, $\underline{v} = (v_1, \ldots, v_s) \in G$, let

$$J^{+I}(\underline{v}) := \{g \in J(\underline{v}) : v_i(g) > v_i \text{ for } i \in I\},$$

$J^+(\underline{v}) := J^{+I_0}(\underline{v})$.

Definition: The Poincaré series $P_{\{v_i\}}(t_1, \ldots, t_s)$ of the collection of valuations $\{v_i\}$ is the element of the ring $\mathbb{Z}[[S]]$ defined by

$$P_{\{v_i\}}(\underline{t}) = \sum_{\underline{v} \in S} \left(\sum_{I \subset I_0} (-1)^{\#I} \dim \left(J^{+I}(\underline{v}) / J^{+}(\underline{v}) \right) \right) \cdot \underline{t}^{\underline{v}}.$$

One can see that, for collections of integer valued valuations, this definition coincides with that used e.g. in [1]. One can easily extract a proof of this from the proof of Theorem 3 in [1].

Remark. This notion is defined if all the factor spaces $J^{+I}(\underline{v})/J^{+}(\underline{v})$ have finite dimensions. This takes place if each valuation v_i, $i = 1, \ldots, s$, is centred at the origin, i.e. $\{g \in \mathcal{O}_{X,0} : v_i(g) > 0\}$ coincides with the maximal ideal $\mathfrak{m}$ of the ring $\mathcal{O}_{X,0}$.

Definition: A valuation v with values in a group G is finitely determined if, for each $v_0 \in G$, the condition $v(g) = v_0$ is a constructible condition on a jet of the germ g of a certain (finite) order (see [2]).

Examples. **1.** For an irreducible plane curve germ $C = \{f = 0\} \subset (\mathbb{C}^2, 0)$, let $\varphi : (\mathbb{C}, 0) \to (\mathbb{C}^2, 0)$ be a parametrization (uniformization) of the curve C. For a germ $g \in \mathcal{O}_{X,0}$, let $v_C(g)$ be the order of zero of the function $g \circ \varphi(\tau)$ at the origin (if $g \circ \varphi \equiv 0$, $v_C(g) := +\infty$). The map $v_C : \mathcal{O}_{X,0} \to \mathbb{Z} \cup \{+\infty\}$ is a finitely determined rank 1 valuation on the ring $\mathcal{O}_{X,0}$. (This valuation has a non-trivial kernel, i.e. $v_C(g) = +\infty$ for some $g \neq 0$.)
2. For the same curve $C = \{f = 0\}$ and for $g \in \mathcal{O}_{X,0}$, $g \neq 0$, let $g = f^{k_C(g)} g'$, where g' is not divisible by f. The map $w_C : \mathcal{O}_{X,0} \to \mathbb{Z}^2 \cup \{+\infty\}$, $w_C(g) := (k_C(g), v_C(g'))$, is a rank 2 valuation on the ring $\mathcal{O}_{X,0}$. (The group $\mathbb{Z}^2$ is ordered lexicographically, i.e. $(k', v') > (k'', v'')$ iff $k' > k''$ or $k' = k''$ and $v' > v''$.) The valuation w_C is not finitely determined: for $k > 0$, the condition $k_C(g) = k$ is not determined by a (finite) jet of the germ g. (The kernel of the valuation w_C is trivial.)

The notion of integration with respect to the Euler characteristic χ over the projectivization $\mathbb{P}\mathcal{O}_{X,0}$ of the ring $\mathcal{O}_{X,0}$ was defined in [2]. The argument from [2] give the following statement.

Proposition 1. *If all the valuations v_i, $i = 1, \ldots, s$, are finitely determined, one has*

$$P_{\{v_i\}}(\underline{t}) = \int_{\mathbb{P}\mathcal{O}_{X,0}} \underline{t}^{\,\underline{v}(g)} d\chi \,. \tag{1}$$

Remark. One can formulate the notion of the Poincaré series of a collection of valuations in terms of the extended semigroup of the collection in spirit of [1], [6].

2 Valuations on $\mathcal{O}_{\mathbb{C}^2,0}$

Let $\pi : (X, D) \to (\mathbb{C}^2, 0)$ be a modification of the plane by a (finite) sequence of blowing-ups. The exceptional divisor D is the union of irreducible components E_σ ($\sigma \in \Gamma$); each of them is isomorphic to the complex projective line $\mathbb{CP}^1$. The dual graph of the modification π consists of vertices corresponding to the irreducible components E_σ, i.e. to elements of Γ; two vertices are connected by an edge iff the corresponding components intersect. To $\sigma \in \Gamma$, i.e. to an irreducible component E_σ of the exceptional divisor, there corresponds a natural valuation v_σ: the *divisorial* one. For a germ $g \in \mathcal{O}_{X,0}$ the value $v_\sigma(g)$ is defined as a multiple of (say, c_σ times) the order $w_\sigma(g)$ of zero of the function $g \circ \pi$ along the component E_σ (i.e. of the coefficient at $[E_\sigma]$ in the zero divisor of the function $g \circ \pi$). It is convenient to choose the coefficient c_σ in such a way that $\min\limits_{g\in\mathfrak{m}} v_\sigma(g) = 1$, i.e. $c_\sigma = 1/w_\sigma(g)$ for a generic function $g \in \mathfrak{m}$.

It is known ([8], see also [5]) that all valuations on the ring $\mathcal{O}_{X,0}$ centred at the origin (i.e. such that $v(g) > 0$ for $g \in \mathfrak{m}$) are given by the following list.

I. All rank 1 valuations centred at the origin correspond to some sequences (finite or infinite) of blowing-ups such that each next blowing-up is made at a point of the divisor born on the previous step. To get a one-to-one correspondence, one should exclude sequences of blowing-ups made at each step after a certain one at the intersection point of a fixed divisor E_{σ_0} with the last one. (In such case the correspondence described below leads to the divisorial valuation v_{σ_0} associated to the divisor E_{σ_0}.) If the sequence is finite, the corresponding valuation is the divisorial one associated to the last divisor. If the sequence is infinite, the value $v(g)$ of the corresponding valuation is defined as the limit $\lim\limits_{i\to\infty} v_{\sigma_i}(g)$ where v_{σ_i} is the divisorial valuation associated to the divisor born on i-th step. Depending on the sequence of blowing-ups, one can distinguish the following types of valuations.

I.1. The blowing-ups are made at the intersection points of the strict transforms of a fixed irreducible curve $C = \{f = 0\} \subset (\mathbb{C}^2, 0)$ with the exceptional divisor. The corresponding valuation (a *curve valuation of rank 1*) is equivalent to the valuation defined by the order of a function g on the curve C in an uniformization parameter (Example 1 above). The corresponding dual graph of the modification has a growing infinite tail: Fig.1. This valuation has a non-trivial kernel.



Figure 1: The dual graph of a valuation of type I.1.

I.2. The sequence of blowing-ups produces an infinite tail like on Fig.1, but there exists no curve corresponding to this sequence (one can say that there is a formal curve defined by a formal power series), one gets a discrete valuation with the trivial kernel: a *formal curve valuation*.

I.3. The sequence of blowing-ups increases the number of rupture points producing, as the limit, the graph shown on Fig.2. In this case the group of values is contained in the ring $\mathbb{Q}$ of rational numbers and is not finitely generated. For any $g \in \mathcal{O}_{\mathbb{C}^2,0}$, one has $v(g) = v_{\sigma_i}(g)$ for i sufficiently large. Valuations of this sort we call *infinite valuations*.



Figure 2: The dual graph of a valuation of type I.3.

I.4. In the cases I.2 and I.3 blowing-ups were made either permanently after a certain step or from time to time at a smooth point of the last divisor. If this is not the case (i.e. if all the blowing-ups starting from a moment are produced at an intersection point of the last divisor with a previous one but not with one and the same) one gets a valuation with a finitely generated group of values, which is not contained in $\mathbb{Q}$ (an *irrational valuation*).

II. Besides that there are two kinds of rank 2 valuations on the ring $\mathcal{O}_{\mathbb{C}^2,0}$.

II.1. For an irreducible curve germ $C = \{f = 0\} \subset (\mathbb{C}^2, 0)$ one has the valuation described in Example 2: *curve valuations of rank 2*. Valuations of this type are the only ones on the list which are not finitely determined.

II.2. For a divisor E_σ of a modification $\pi : (X, D) \to (\mathbb{C}^2, 0)$ and for a point P on it, let E_σ be locally given by an equation $u = 0$. Then we can define a rank 2 valuation $v = v(g)$ with values in the group $\mathbb{Z}^2$ ordered lexicographically applying the construction of II.1 to the germ of function $g \circ \pi$ at the point P of the curve $u = 0$. This means that one writes $g \circ \pi$ as $u^{v_\sigma(g)} g'$ where $g' \notin \langle u \rangle$ ($v_\sigma(g)$ is just the value of g with respect to the divisorial valuation v_σ) and puts $v(g) := (v_\sigma(g), w_u(g'))$, where $w_u(g')$ is the order of zero of the function g' on the smooth curve $u = 0$ at the point P. This valuation is called an *exceptional curve valuation*.

If one excludes valuations of type I.1. (i.e. those which have a non-trivial kernel), one can say that all valuations on the ring $\mathcal{O}_{\mathbb{C}^2,0}$ are in one to one correspondence with all (finite and infinite) sequences of blowing-ups of the described type. In this case sequences of blowing-ups excluded in I correspond to valuations of type II.2. To a sequence of the type described in I.1 one associates the valuation II.1 (of rank 2) corresponding to the curve germ.

Remarks. **1.** A valuation of type I.1 is essentially the second component of the corresponding valuation of type II.1. If one considers the ring $\mathbb{C}[[x, y]]$ of formal power series in two variables instead of $\mathcal{O}_{\mathbb{C}^2,0}$, valuations of type I.1 and I.2 constitute one and the same type. The results of the paper are valid in this setting as well.

2. Except valuations centred at the origin, there are so called f-adic valuations k_C corresponding to irreducible plane curve singularities $C = \{f = 0\} \subset (\mathbb{C}^2, 0)$. For a germ $g \in \mathcal{O}_{\mathbb{C}^2,0} \setminus \{0\}$, $k_C(g)$ is defined by the relation $g = f^{k_C(g)} g'$ where $g' \notin \langle f \rangle$. This is just the first component of the rank 2 curve valuation corresponding to C. For an appropriate definition, the Poincaré series of a collection of valuations containing f-adic ones is a reduction of the Poincaré series of the collection obtained by substituting f-adic ones by the corresponding plane curve valuations of rank 2.

For further discussions it is convenient to use a notion of a resolution of a (finite) collection of valuations. For a divisorial valuation this is a modification $\pi : (X, D) \to (\mathbb{C}^2, 0)$ of the plane by a finite sequence of blowing-ups over the origin which contains the corresponding divisor (this means that if is a modification of the minimal resolution by a finite sequence of blowing-ups). For a valuation of one of the types I, II.1, let $\pi_i : (X_i, D_i) \to (\mathbb{C}^2, 0)$ be the modification obtained at the i-th step of the corresponding sequence of blowing-ups. Let (X, D) be the projective limit (in the category of analytical spaces) of the sequence $\{(X_i, D_i)\}$ and let $\pi : (X, D) \to (\mathbb{C}^2, 0)$ be the corresponding map. One can see that X is a smooth complex surface and D is the union of infinitely many projective lines on it. The map $\pi : (X, D) \to (\mathbb{C}^2, 0)$ will be called the minimal resolution of the valuation. This is not a resolution since since, in particular, the map π is not proper. For a valuation of type II.2, the minimal resolution will be defined as the minimal resolution of the corresponding divisorial valuation (the first component of the considered one) followed by an additional blowing-up at the corresponding point. Thus in the minimal resolution the point under consideration is an intersection point of components of the exceptional divisor.

The minimal resolution of a finite collection of valuations is the projective limit of the corresponding (multi-index) system of modifications. It is simply the fibre product of the minimal resolutions of all valuations.

Finally a resolution of a finite collection of valuations is a modification of the space (X, D) of the minimal resolution by a finite number of blowing-ups (at points of D).

3 Poincaré series of a collection of plane valuations

Let v_i, $i = 1, \ldots, s$, be a set of valuations on the ring $\mathcal{O}_{\mathbb{C}^2,0}$ centred at the origin. Let $\pi : (X, D) \to (\mathbb{C}^2, 0)$ be a resolution of the set of valuations $\{v_i\}$. For a component E_σ of the exceptional divisor D, $\sigma \in \Gamma$, let L_σ be a smooth germ of curve transversal to D at a smooth point of it, let the curve germ $\ell_\sigma = \pi(L_\sigma) \subset (\mathbb{C}^2, 0)$ be given by an equation $\{g_\sigma = 0\}$, and let $m_i^\sigma = v_i(g_\sigma)$, $\underline{m}^\sigma := (m_1^\sigma, \ldots, m_s^\sigma) \in G$.

Without loss of generality one can suppose that $v_1, \ldots, v_r$ are valuations of type II.1 (or rank 2) and the others are not. For a component E_σ of the exceptional divisor D, let $\overset{\circ}{E}_\sigma$ be its "smooth part", i.e. E_σ itself minus the intersection points with all other components of D. For $i = 1, \ldots, r$, let $C_i = \{f_i = 0\}$ be the corresponding curve, $f_i \in \mathcal{O}_{\mathbb{C}^2,0}$.

Theorem 1. *The Poincaré series of the set of valuations $\{v_i\}$ is given by the* *equation*

$$P_{\{v_i\}}(\underline{t}) = \prod_{\sigma\in\Gamma}(1-\underline{t}^{\,\underline{m}^\sigma})^{-\chi(\overset{\circ}{E}_\sigma)} \times \prod_{i=1}^{r}(1-t_i^{(1,0)}\prod_{j\neq i} t_j^{v_j(f_i)})^{-1}\,. \tag{2}$$

Proof. One cannot use the equation (1) directly since valuations of type II.1 are not finitely determined. However the condition $v_i(g) = (0, v)$ is defined by a jet of the germ g of a finite order. Therefore let us first compute the part $P^o(\underline{t})$ of the Poincaré series $P_{\{v_i\}}(\underline{t})$ which consists of monomials which are not divisible by any of $t_i^{(1,0)}$, $i = 1, \ldots, r$. One can see that

$$P^o(\underline{t}) = \int\limits_{\mathbb{P}(\mathcal{O}_{\mathbb{C}^2,0}\setminus\bigcup\limits_{i=1}^{r}\langle f_i\rangle)} \underline{t}^{\,\underline{v}} d\chi$$

Computation of $P^o(\underline{t})$ essentially repeats the arguments from, e.g., [2] or [6]. To compute the series $P^o(\underline{t})$ up to terms of any fixed degree $\underline{V} \in G$ one can make finitely many additional blowing-ups at intersection points of the components of the exceptional divisor D so that, for a germ $g \in \mathcal{O}_{\mathbb{C}^2,0} \setminus \bigcup\limits_{i=1}^{r}\langle f_i\rangle$ with $\underline{v}(g) \leq \underline{V}$, all the intersection points of the strict transform of the curve $\{g = 0\}$ with the exceptional divisor D belong to its smooth part $\overset{\circ}{D} = \bigcup\limits_{\sigma} \overset{\circ}{E}_\sigma$. Let $\mathcal{O}^{\underline{V}}_{\mathbb{C}^2,0} = \{g \notin \bigcup\limits_{i=1}^{r}\langle f_i\rangle : \underline{v}(g) \leq \underline{V}\}$, and let $\mathcal{O}^{D}_{\mathbb{C}^2,0}$ be the set of germs $g \in \mathcal{O}_{\mathbb{C}^2,0} \setminus \{0\}$ such that the intersection points of the strict transform of the curve $\{g = 0\}$ with the exceptional divisor D belong to $\overset{\circ}{D}$. The integral

$$\int_{\mathbb{P}\mathcal{O}^{D}_{\mathbb{C}^2,0}} \underline{t}^{\,\underline{v}} d\chi$$

over the projectivization $\mathbb{P}\mathcal{O}^{D}_{\mathbb{C}^2,0}$ contains all terms of the series $P^o(\underline{t})$ up to degree $\underline{V}$. There is a map from $\mathbb{P}\mathcal{O}^{D}_{\mathbb{C}^2,0}$ to the space of effective divisors on $\overset{\circ}{D}$: to a function germ $g \in \mathcal{O}^{D}_{\mathbb{C}^2,0}$ one associates the intersection of the strict transform of the curve $\{g = 0\}$ with the exceptional divisor D. Proposition 2 from [2] implies that the preimage of a point with respect to this map is a complex affine space and thus has the Euler characteristic equal to 1. The Fubini formula implies that the integral $\int_{\mathbb{P}\mathcal{O}^{D}_{\mathbb{C}^2,0}} \underline{t}^{\,\underline{v}} d\chi$ is equal to the integral with respect to the Euler characteristic of the monomial $\underline{t}^{\,\underline{v}}$ over the space of effective divisors on $\overset{\circ}{D}$, where $\underline{v}$ is the additive function on the space of effective divisors (with values in G) equal to $\underline{m}^\sigma$ for a point from the component $\overset{\circ}{E}_\sigma$ of $\overset{\circ}{D}$.

The space of effective divisors on $\overset{\circ}{D}$ is the direct product of the spaces of effective divisors on the components $\overset{\circ}{E}_\sigma$, $\sigma \in \Gamma$. Each of the latter ones is the disjoint union of the symmetric powers $S^k\ \overset{\circ}{E}_\sigma$ of the component $\overset{\circ}{E}_\sigma$. Therefore

$$\int_{\mathbb{P}\mathcal{O}^D_{\mathbb{C}^2,0}} \underline{t}^{\underline{v}} d\chi = \prod_{\sigma\in\Gamma}\left(\sum_{k=0}^{\infty}\chi(S^k\ \overset{\circ}{E}_\sigma)\cdot \underline{t}^{k\underline{m}^\sigma}\right)\ .$$

Using the equation

$$\sum_{k=0}^{\infty}\chi(S^kX)\cdot t^k = (1-t)^{\chi(X)}$$

one gets

$$\int_{\mathbb{P}\mathcal{O}^D_{\mathbb{C}^2,0}} \underline{t}^{\underline{v}}\ d\chi = \prod_{\sigma}(1-\underline{t}^{\underline{m}^\sigma})^{-\chi(\overset{\circ}{E}_\sigma)}\ . \tag{3}$$

The right hand side of equation 3 do not contain components of the exceptional divisor born under additional blowing-ups since for each of them $\chi(\overset{\circ}{E}_\sigma)=0$. Therefore

$$P^o(\underline{t}) = \prod_{\sigma\in\Gamma}(1-\underline{t}^{\underline{m}^\sigma})^{-\chi(\overset{\circ}{E}_\sigma)}$$

(for a resolution $\pi:(X,D)\to(\mathbb{C}^2,0)$ of the set of valuations $\{v_i\}$).

Now, for $\underline{k}=(k_1,\ldots,k_r)\in\mathbb{Z}^r_{\geq 0}$, let $P^{\underline{k}}(\underline{t})$ be the sum of terms of the Poincaré series $P_{\{v_i\}}(\underline{t})$ divisible by $\prod\limits_{i=1}^{r} t_i^{(k_i,0)}$ but not divisible by a monomial of this sort of higher degree. The set of functions g with $\underline{v}(g)$ divisible by $\prod\limits_{i=1}^{r} t_i^{(k_i,0)}$ (but not by a monomial of higher degree) is just $\prod\limits_{i=1}^{r} f_i^{k_i}\cdot\left(\mathcal{O}_{\mathbb{C}^2,0}\setminus\bigcup\limits_{i=1}^{r}\langle f_i\rangle\right)$. One has $v_j(\prod\limits_{i=1}^{r} f_i^{k_i})=\sum\limits_{i=1}^{r}k_iv_j(f_i)$ and $v_i(f_i)=(1,0)$ for $i=1,\ldots,r$.

From this it follows that

$$P^{\underline{k}}(\underline{t}) = P^o(\underline{t})\prod_{i=1}^{r}t_i^{(k_i,0)}\prod_{j\neq i}t_j^{k_iv_j(f_i)}$$

and therefore

$$P_{\{v_i\}}(\underline{t}) = P^o(\underline{t})\sum_{\underline{k}\in\mathbb{Z}^r_{\geq 0}}\prod_{i=1}^{r}t_i^{(k_i,0)}\prod_{j\neq i}t_j^{k_iv_j(f_i)} = P^o(\underline{t})\cdot\prod_{i=1}^{r}(1-t_i^{(1,0)}\prod_{j\neq i}t_j^{v_j(f_i)})^{-1}\ .$$

□

4 Poincaré series determines dual graphs

To a resolution $\pi:(X,D)\to(\mathbb{C}^2,0)$ of a collection of valuations one associates the dual graph (generally speaking infinite). It consists of vertices corresponding to the irreducible components of the exceptional divisor D; two vertices are connected by an edge if the corresponding components intersect. The set of vertices of the dual graph inherits a partial order defined by approximation of the modification by sequences of blowing-ups: a component $E_{\sigma'}$ is "greater" than another component E_σ ($\sigma' > \sigma$) if the exceptional divisor of the minimal modification which contains $E_{\sigma'}$ also contains E_σ. Two resolutions are called combinatorially equivalent if their dual graphs together with the partial orders are isomorphic.

For collections of valuations of types I.1 and I.2 the Poincaré series coincides with the Alexander polynomial (in several variables) of the corresponding algebraic link (obtained by cutting the non-convergent series for a valuation of type I.2) (see [1]). It is known that the Alexander polynomial of an algebraic link determines the topology of the curve singularity ([7], see another proof in [3]). Therefore the Poincaré series of a collection of such valuations determines its minimal resolution up to combinatorial equivalence.

In [3], it was shown that the Poincaré series of a collection of divisorial valuations "determines the topology of the set of divisors" in the sense that it determines the dual graph of the minimal resolution up to combinatorial equivalence. Moreover, it was shown that the Poincaré series of a collection of valuations which includes both divisorial ones and those of types I.1 (or I.2) does not determine, in general, the dual graph of the minimal resolution.

Theorem 2. *Suppose that a collection $\{v_i\}$ of valuations does not contain valuations if types I.1 and I.2. Then the Poincaré series $P_{\{v_i\}}(\underline{t})$ of the collection determines types of the valuations, the dual graph of the minimal resolution up to combinatorial equivalence and divisors or sequences of divisors corresponding to the valuations.*

In what follows, for short, we shall simply say that the Poincaré series determines the dual graph of the minimal resolution.

Proof. One property of collections of valuations of the type under consideration used in the proof is the fact that the projection formula holds for them. This means that, for a subset $I \subset I_0 = \{1, \ldots, s\}$, the Poincaré series of the set of valuations $\{v_i\}_{i \in I}$ is obtained from the Poincaré series of the whole set $\{v_i\}$ by omitting the variables t_i with $i \notin I$ (in other words by substituting t_i with $i \notin I$ by 1).

Remark. Omitting a valuation of type I.3 (an infinite valuation), as a factor one formally gets an infinite product of the form

$$(1 - \underline{t}^{\underline{m}^1})(1 - \underline{t}^{\underline{m}^1})^{-1}(1 - \underline{t}^{\underline{m}^2})(1 - \underline{t}^{\underline{m}^2})^{-1} \ldots$$

This product should be canceled.

The projection formula follows directly from the equation (2) (Theorem 1).

Remark. For collections of valuations which do not contain valuations of type II.1 (curve valuations of rank 2) this can be also deduced from equation (1). The fact that the projection formula holds for valuations of type II.1 as well can mean that this valuations can be considered finitely determined in some weak sense.

The projection formula implies that, from the Poincaré series of a collection of valuations, one can restore, in particular, the Poincaré series of each individual valuation from the collection.

For each individual valuation one can define its type from the Poincaré series. From equation (2) it follows that the set of exponents in the Poincaré series with non-vanishing coefficients generates an abelian group of rank equal to the rank of the valuation. The Poincaré series of a valuation of type II.1 (a curve valuation of rank 2) has infinitely many non-vanishing terms with exponents from the isolated subgroup of rank 1. This does not take place for valuations of type II.2 (exceptional curve valuations).

Each series from the ring $Z[[S]]$ with the free term equal to 1 has a unique representation of the form $\prod\limits_{a\in S}(1-t^a)^{k_a}$ with $k_a \in \mathbb{Z}$. (Generally speaking, this product is not finite: infinite number of the exponents k_a may be different from zero.)

The Poincaré series of rank 1 valuations have the following form.

1) For a divisorial valuation

$$P(t) = \frac{\prod\limits_{i=1}^{h}(1-t^{m^{\alpha_i}})}{\prod\limits_{i=0}^{h+1}(1-t^{m^{\beta_i}})} \tag{4}$$

where the exponents m^{β_i} generate an infinite cyclic group (i.e. a group isomorphic to $\mathbb{Z}$). Here α_i are rupture points and β_i are dead ends of the dual graph of the minimal resolution.

2) For a valuation of type II.4 (an irrational one) the Poincaré series has the same form 4, but the exponents m^{β_i} generate a free abelian group of rank 2. (The ratios m^{β_i}/m^{β_0} are rational for $i \leq h$ and the ratio $m^{\beta_{h+1}}/m^{\beta_0}$ is irrational.

3) For a valuation of type II.3 (an infinite valuation) the Poincaré series has a representation of the form

$$P(t) = \frac{\prod\limits_{i=1}^{\infty}(1-t^{m^{\alpha_i}})}{\prod\limits_{i=0}^{\infty}(1-t^{m^{\beta_i}})}$$

with infinitely many factors.

This shows that rank 1 valuations of different types cannot have equal Poincaré series.

Remark. This also can be deduced from the fact that, for one valuation, the set of exponents with non-zero coefficients coincides with the semigroup of values of the valuation.

For a collection of divisorial valuations the statement of the Theorem was proved in [3]. We shall reduce consideration of a collection of valuations of different types to the case of a collection of divisorial valuations. For that we shall substitute each non-divisorial valuation from the collection by one or two divisorial ones. In each case the Poincaré series of the resulting collection of valuations should be defined by the Poincaré series of the initial collection and the dual graph of the minimal resolution of the resulting collection (or rather os series of them) should permit to restore the dual graph of the minimal resolution of the initial one.

Suppose that a valuation from the collection $\{v_i\}$ is of type II.1 (a curve valuation of rank 2). Without loss of generality we may assume that this is the first one. Let E_{σ_0} be a vertex of the dual graph of the minimal resolution of the collection $\{v_i\}$ far enough on the infinite tail corresponding to the valuation v_1. Let us substitute the (rank 2) valuation v_1 in the collection $\{v_i\}$ by the divisorial valuation v_{σ_0}. If one knows the dual graph of the minimal resolution of the new collection, one can easily restore the dual graph for the initial one. One can see that the Poincaré series of the new collection of valuations is obtained from the Poincaré series $P_{\{v_i\}}(t_1, \ldots, t_s)$ (see equation (2)) by substituting $t_1^{(0,1)}$ by t_{σ_0} and $t_1^{(1,0)}$ by $t_{\sigma_0}^N$ with N large enough.

Suppose that the valuation v_1 is of type II.2 (an exceptional curve valuation), let E_{σ_0} be the exceptional curve (a component of the exceptional divisor) corresponding to the valuation, and let P be the corresponding point of the component E_{σ_0}. (The point P is the intersection point of the component E_{σ_0} with another component of the exceptional divisor.) Let us make sufficiently many additional blowing-ups at the point P of the component E_{σ_0}. (If one knows the dual graph of a resolution obtained this way, one can easily restore the dual graph for the minimal one.) In the new resolution, let P be the intersection point of the component E_{σ_0} with a component $E_{\sigma_0'}$. Let us substitute, in the collection $\{v_i\}$, the valuation v_1 by two divisorial valuations v_{σ_0} and $v_{\sigma_0'}$. One can see that the Poincaré series of the new collection of valuations is obtained from the Poincaré series $P_{\{v_i\}}(t_1, \ldots, t_s)$ by substituting $t_1^{(1,0)}$ by t_{σ_0} and $t_1^{(0,1)}$ by $t_{\sigma_0'} t_{\sigma_0}^N$ with N large enough.

Suppose that v_1 is a valuation of type I.3 (an infinite one). Moreover let us assume that $v_1, v_2, \ldots, v_r$ are all valuations of type I.3 in the collection. The Poincaré series has the form

$$\prod_{n=1}^{n_0}(1-\underline{t}^{\underline{k}^n})^{-\chi_n}\prod_{\ell=1}^{r}\prod_{j=1}^{\infty}\frac{1-\underline{t}^{\underline{m}^{\alpha_j^\ell}}}{1-\underline{t}^{\underline{m}^{\beta_j^\ell}}}\,.$$

Here the product $\prod\limits_{j=1}^{\infty}\frac{1-\underline{t}^{\underline{m}^{\alpha_j^\ell}}}{1-\underline{t}^{\underline{m}^{\beta_j^\ell}}}$ corresponds to the infinite part (tail) of the dual graph corresponding to the valuation v_ℓ. In these products one has

$$\underline{m}^{\alpha_1^\ell} < \underline{m}^{\beta_1^\ell} < \underline{m}^{\alpha_2^\ell} < \underline{m}^{\beta_2^\ell} < \ldots$$

and $\underline{m}^{\alpha_j^\ell} \to \infty$, $\underline{m}^{\beta_j^\ell} \to \infty$ when $j \to \infty$. The distribution of the factors between these products (from a certain place) is determined by the following property: the ratio $m_{j_1}^{\beta_j^\ell}/m_{j_2}^{\beta_j^\ell}$ strictly increases with j along the tail corresponding to the valuation v_{j_1} (i.e. for $\ell = j_1$), strictly decreases along the tail corresponding to the valuation v_{j_2} ($\ell = j_2$) and is constant along all other tails.

Let us substitute the infinite valuation v_1 by the divisorial valuation $v_{\alpha_N^1}$ with N large enough. The dual graph of the minimal resolution of the collections $\{v_{\alpha_N^1}, v_2, \ldots, v_s\}$ is obtained from the dual graph of the minimal resolution of the collections $\{v_i\}$ by truncation of the infinite tail corresponding to the valuation v_1 at the vertex α_N^1. The dual graph of the resolution of the collection $\{v_i\}$ can be restored if one knows the truncated ones for all N large enough. The Poincaré series of the collection $\{v_{\alpha_N^1}, v_2, \ldots, v_s\}$ is equal to

$$\prod_{n=1}^{n_0}(1-\underline{t}^{\underline{k}^n})^{-\chi_n}\prod_{j=1}^{N}\frac{1-\underline{t}^{\underline{m}^{\alpha_j^1}}}{1-\underline{t}^{\underline{m}^{\beta_j^1}}}\prod_{\ell=2}^{r}\prod_{j=1}^{\infty}\frac{1-\underline{t}^{\underline{m}^{\alpha_j^\ell}}}{1-\underline{t}^{\underline{m}^{\beta_j^\ell}}}\,.$$

Suppose that the valuation v_1 is of type II.4 (an irrational one). The semigroup of values of the valuation v_1 is generated by $1 = m_1^{\beta_0}, m_1^{\beta_1}, m_1^{\beta_2}, \ldots, m_1^{\beta_h}$ where $m_1^{\beta_0} < m_1^{\beta_1} < m_1^{\beta_2} < \ldots < m_1^{\beta_h}$, the numbers $m_1^{\beta_i}$ are rational for $i < h$ and $m_1^{\beta_h}$ is irrational. Let us substitute the valuation v_1 in the collections $\{v_i\}$ by the divisorial valuation v_{σ_h} far enough in the sequence $\{\sigma_h\}$ corresponding to the valuation v_1 and moreover such that in the dual graph of the minimal resolution of the valuation v_1 the component E_{σ_h} does not intersect the component $E_{\sigma_{h+1}}$ (the first among those born by blowing-ups at points of the component E_{σ_h}). The dual graph of the resolution of the collection $\{v_i\}$ can be restored if one knows the dual graphs of the minimal resolutions of the collections $\{v_{\sigma_h}, v_2, \ldots, v_s\}$ with all h large enough of the described type. The Poincaré series of the collection $\{v_{\sigma_h}, v_2, \ldots, v_s\}$ is obtained from the Poincaré series $P_{\{v_i\}}(\underline{t})$ of the collection $\{v_i\}$ by substituting each monomial $t_1^{m_1}t_2^{m_2}\ldots t_s^{m_s}$ by $t_1^{m_1'}t_2^{m_2}\ldots t_s^{m_s}$ where m_1' is defined in the following way. Let α_N be the result of the truncation of the continuous fraction of the (irrational) number $m_1^{\beta_h}$ at the level N large enough. Let $m_1 = k_0 m_1^{\beta_0} + \cdots + k_{h-1}m_1^{\beta_{h-1}} + k_h m_1^{\beta_h}$. Then $m_1' = k_0 m_1^{\beta_0} + \cdots + k_{h-1}m_1^{\beta_{h-1}} + k_h\alpha_N$. □

References

[1] Campillo A., Delgado F., Gusein-Zade S.M. The Alexander polynomial of a plane curve singularity via the ring of functions on it. Duke Math. J., v.117, no.1, 125–156 (2003).

[2] Campillo A., Delgado F., Gusein-Zade S.M. The Alexander polynomial of a plane curve singularity and integrals with respect to the Euler characteristic. International Journal of Maths., v.14, no.1, 47–52 (2003).

[3] Campillo A., Delgado F., Gusein-Zade S.M. The Poincaré series of divisorial valuations in the plane defines the topology of the set of divisors. To appear in Functional Analysis and Other Mathematics. arXiv:0806.4492. (2008).

[4] Campillo A., Galindo, C. On the graded ring relative to a valuation. Manuscr. Math., v.92. 173–189 (1997).

[5] Delgado F., Galindo C., Nuñez A. Saturation for valuations on two-dimensional regular local rings. Math. Zeitschrift., v.234, no.3, 519–550 (2000).

[6] Delgado F., Gusein-Zade S.M. Poincaré series for several plane divisorial valuations. Proc. Edinb. Math. Soc. (2), v.46, no.2, 501–509 (2003).

[7] Yamamoto M. Classification of isolated algebraic singularities by their Alexander polynomials. Topology, v.23, no.3, 277–287 (1984).

[8] Zariski O., Samuel, P. Conmutative Algebra. Vol. II. Van Nostrand. Princeton N.J. (1960).